2365

Psyche Vol. 59 Publ. August 1952

other. *F. ulkei* is a boreal form whose main range is in the northern tier of states and southern Canada.

LITERATURE CITED

CREIGHTON, W. S.
1950. The ants of North America. Bull Mus. Comp. Zool., 104:1-585.

EMERY, C.
1893. Beiträge zur Kenntnis der nordamerikanischen Ameisen-fauna. Zool. Jahrb. Syst., 7:633-682.

WHEELER, W. M.
1913. A revision of the ants of the genus *Formica*. Bull. Mus. Comp. Zool., 53:379-565.

CORRECTION TO THE SYNONYMY OF THE ANT CAMPONOTUS FORMOSENSIS WHEELER.—In a joint paper (Yasumatsu and Brown, 1951, Jour. Fac. Agr., Univ. Kyushu, 10, cf. p. 42), the synonymy of *Camponotus formosensis* is cited incorrectly. Due to an unfortunate clerical error, the name *Camponotus maculatus taylori* var. *formosae* Wheeler and the variant nomenclatorial equivalents *C. barbatus albosparsa* var. *formosae*, Emery and *C. barbatus taylori* var. *formosae*, Teranishi were wrongly included in this formal list of synonyms and must now be removed. The variety *formosae*, whatever may be its relationship to *maculatus* or *barbatus*, is of course not at all closely related to *C. formosensis* or any other member of the *C. herculeanus* group.—W. L. BROWN, JR., Museum of Comparative Zoology, Harvard University.

B